


# Simulation and sensitivities for a phased IceCube-Gen2 deployment

Brian Clark, Rob Halliday  
*for the IceCube-Gen2 Collaboration*  
Michigan State University



ICRC 2021  
“Berlin”



MICHIGAN STATE  
UNIVERSITY

# Questions

- What we know about the flux of high-energy neutrinos
  - Roughly power law in shape
  - Seemingly flavor democratic
  - Consistent with isotropic arrival direction
- But...
  - No definitive sources (some tantalizing evidence!)
  - No UHE neutrinos
- Goals for a next generation instrument:
  - Better pointing for point source localization and multi-messenger observations → improved angular resolution
  - Statistically significant observations over a broad energy range




# IceCube-Gen2


A broadband neutrino observatory

Four new elements, leveraging complimentary technologies, to achieve sensitivity to MeV-EeV neutrinos

1. IceCube Upgrade
2. Enlarged deep optical array
3. Surface Array
4. Shallow radio array




# IceCube-Gen2



# Gen2-Optical

## TeV-EeV neutrinos

- 8 km<sup>3</sup> optical array: ~10x volume of IceCube
- Laid out in a “Sunflower” pattern
  - 120 strings, 240m lateral spacing
  - 80 OMs/string, 17m vertical spacing
- Larger depth range than IceCube
  - Gen2: 1340-2700 m
  - IceCube: 1446-2451 m
- Deployment expected to take ~7 seasons, with ~21 holes drilled per season
  - Drill rate ramps up in first two seasons




# Gen2-Optical

## Event Topologies

- Like IceCube (and many other telescopes!) two primary detection channels: cascades and tracks
- Focus on *throughgoing* tracks
  - $\nu_\mu/\bar{\nu}_\mu$  charged-current interactions
  - *Vertex outside* of the contained volume
- This class of events provides the longest lever arm for reconstruction of neutrino direction
- Provides best sensitivity in search for sources of neutrinos


### Cascades

$\nu_e/\bar{\nu}_e, \nu_\tau/\bar{\nu}_\tau$  charged current  
All flavors neutral current




### Tracks

Mostly  $\nu_\mu/\bar{\nu}_\mu$  charged current




# Simulation

- Hard, isotropic flux of single high energy muons (3 TeV-100 PeV,  $E^{-1.4}$ )
- Photons are propagated in ice (*CLsim* code)
  - Photons weighted by module wavelength acceptance for efficiency
- Received photons are processed into recorded hits
- Models of noise and triggers are applied
- Finally, apply likelihood based reconstruction





IceCube-Gen2 (B. A. Clark)



# Performance

- Apply quality cuts to isolate well-reco'd tracks
  - Minimum number of hit OMs
  - Minimum reco track length, etc.
- Two key figures of merit
  - Muon effective area
  - Median angular error (“angular resolution”)
- Halfway through construction, IceCube-Gen2 will already have:
  - 2x the aperture for horizontal events
  - 50% better angular resolution






# Sensitivity

## Steady Sources

- Estimated 1-year discovery potential for  $E^{-2}$  spectrum
- Background assumed to be conventional + prompt atmospheric neutrino flux (Honda, Enberg, + Gaisser H3a)
- Flattening at  $\sin \delta < -0.5$  due to growing surface veto

1 yr Discovery Potential,  $E^{-2.0}$




Season	# Deployed Strings
1	19
3	51
5	87

# Sensitivity


## Flaring Sources

- Estimated significance of detection of a flare as a function of flare duration
- Flare model: 2014/15 flare of TXS0506+056
  - $E^{-\gamma}$ , with  $\gamma = 2.2 \pm 0.2$
  - 156 days long in “box shaped” analysis
- 14/15 flare would have been seen in Gen2 at  $> 5 \sigma$  within the first season after deployment



# Conclusions

- Gen2 will be a broadband neutrino observatory with unprecedented capabilities
- An enlarged optical array will be built over ~7 Antarctic seasons
- Even during construction, Gen2 will have rapidly increasing discovery potential



# Backup

# Gen2-Optical Performance

## Through-going tracks

5x the effective area of IceCube  
2x improvement in angular resolution

