



Improvised Explosive Devices and cosmic rays

A. Vásquez-Ramírez*, M. Ariza-Gómez, M. Carrillo-Moreno, V.G. Baldovino-Medrano, H. Asorey and L.A. Núñez
*presenter e-mail: adriana2168921@uis.edu.co




About **60 countries** and territories are still contaminated with Improvised Explosive Devices (IEDs)

Monitor, L. et al. Concord (2020)



It is possible to use cosmic radiation for the detection of IEDs?

Simulation: interaction between an IED and cosmic radiation



1 Dry soil model


Juárez, M. et al., U. de Alicante (2006)

2 Humid soils models

90 wt.% Dry soil + 10 wt.% water
70 wt.% Dry soil + 30 wt.% water

2 Fertilized soil models

Dry soil + ammonium nitrate
(1 ppm and 2 ppm)



$$\text{ANFO} = 94.3\% + 5.7\%$$



Ammonium nitrate
Diesel oil No.2

Using the LAGO-ARTI framework
(for WCDs simulation response)
Sarmiento-Cano, C. et al., PoS ICRC2019 (2020).

A. contra minas, Tech. Rep. Ejército Nacional (2011)

Results

Particles generated from the interaction of soil models with the Bucaramanga secondary flux of 24 h



Conclusions

The interaction between the **main chemical compounds** of the most commonly **IED found in Colombian soils** with the background flux of cosmic rays at Bucaramanga level **generates particles** that can be **detected**, suggesting a possible IED detection criterion.

The number of **protons** with **0.58 MeV** in **mined soils** is around **237%** greater than protons in **dry soil** model, 2278% in **humid soil** (30wt.%) and 688% for **fertilized soil** (2 ppm).

There is an **excess of protons** around **0.58 MeV** in the presence of the IED


Improvised Explosive Devices and cosmic rays

A. Vásquez-Ramírez*, M. Ariza-Gómez, M. Carrillo-Moreno, V.G. Baldovino-Medrano, H. Asorey and L.A. Núñez

*presenter e-mail: adriana2168921@uis.edu.co

About **60 countries** and territories are still contaminated with Improvised Explosive Devices (IEDs)

Monitor, L. et al., *Concord* (2020)




It is possible
to use cosmic
radiation for
the **detection** of
IEDs?

Improvised Explosive Devices and cosmic rays

A. Vásquez-Ramírez*, M. Ariza-Gómez, M. Carrillo-Moreno, V.G. Baldovino-Medrano, H. Asorey and L.A. Núñez
 *presenter e-mail: adriana2168921@uis.edu.co

Simulation: interaction between an IED and cosmic radiation



$$\text{ANFO} = 94.3\% + 5.7\%$$

Ammonium
nitrate Diesel oil
No.2


It is possible
to use cosmic
radiation for
the **detection** of
IEDs?

Improvised Explosive Devices and cosmic rays

A. Vásquez-Ramírez*, M. Ariza-Gómez, M. Carrillo-Moreno, V.G. Baldovino-Medrano, H. Asorey and L.A. Núñez

*presenter e-mail: adriana2168921@uis.edu.co


Results



There is an excess of protons around 0.58 MeV in the presence of the IED

Improvised Explosive Devices and cosmic rays

A. Vásquez-Ramírez*, M. Ariza-Gómez, M. Carrillo-Moreno, V.G. Baldovino-Medrano, H. Asorey and L.A. Núñez
*presenter e-mail: adriana2168921@uis.edu.co



There is an excess of protons around 0.58 MeV in the presence of the IED

Conclusions

The interaction between the main chemical compounds of the most commonly IED found in Colombian soils with the background flux of cosmic rays at Bucaramanga level generates particles that can be detected, suggesting a possible IED detection criterion.

The number of protons with 0.58 MeV in mined soils is around 237% greater than protons in dry soil model, 2278% in humid soil (30wt.\%) and 688% for fertilized soil (2 ppm).