

Study on the cosmic ray intensity variation using scintillation counters for air shower observation


T.Nonaka¹, A.Oshima², K.Yamazaki² for The Telescope Array Collaboration , for The GRAPES-3 Collaboration

1: Institute for Cosmic Ray Research, University of Tokyo 5-1-5 Kashiwanoha Kashiwa Chiba 277-8582 Japan

2: College of Engineering, Chubu University, Kasugai, Aichi 487-8501, Japan

Telescope Array Detectors

Total detector area for atmospheric muon : 1500 m^2



Three observations of different energies




- TA • $10^{18} \text{ eV} \sim 10^{20} \text{ eV}$
- TALE • $10^{16} \text{ eV} \sim 10^{18.5} \text{ eV}$
- TAx4 • $10^{19.7} \text{ eV} \sim 10^{20} \text{ eV}$



TASD:

- Large area surface → collect scintillation light with WLS fiber
- Thin scinchlator → Low threshold. (~30MeV)
- Sensitive to EM / Mu

Geomagnetic Cutoff : 3GV @vertical



Summary

Using Telescope Array Surface Detector , we obtained data comparable with other observations.

- ✓ Total detector area 1500 m^2
- ✓ Maximum time resolution 1min
- ✓ Count rate $\sim \frac{750 \text{ Hz}}{3 \text{ m}^2}$
- ✓ Trigger threshold : $\sim 20 - 30 \text{ Mev}$ for electron, muon
- ✓ Geomagnetic cut off 3 GV @vertical
- ✓ MedianRigidity: 40GV