


A Data-Driven approach for the measurement of $^{10}\text{Be}/^9\text{Be}$ flux ratio in Cosmic Rays with magnetic spectrometers

F. Nozzoli^{1,2} and C. Cernetti²

¹ INFN-TIFPA Trento Institute for Fundamental Physics and Applications. ² Università degli Studi di Trento, Trento, Italy.

The $^{10}\text{Be}/^9\text{Be}$ flux ratio (thanks to the 2 My lifetime of ^{10}Be) is a radioactive clock providing the measurement of CR residence time in the Galaxy. Existing measurements of $^{10}\text{Be}/^9\text{Be}$ in CR are limited to low energy and affected by large uncertainties, in particular from the Montecarlo simulation. A Data-Driven approach in magnetic spectrometers is presented, as an example it is applied to PAMELA data providing a new measurement in the 0.25-0.85 GeV/n range.

Isotopes of Beryllium in cosmic rays



Be isotopes in CR:


^7Be (E.C. $T_{1/2}=55\text{d}$)
("naked" ^7Be is stable)

^9Be (stable)

^{10}Be (β $T_{1/2}=1.39\text{My}$)
(is a radioactive clock)

Be is scarce in CR not produced by stellar nucleosynthesis
Be is a secondary CR produced by CNO spallation with ISM

PAMELA experiment




ToF:
 $\beta = \frac{v}{c}$

TRK:
 $\frac{RZ}{\beta\gamma} = M$

Calo:
 $\frac{dE}{dX}(\beta)$

Mass & R resolution



$$\frac{\Delta M}{M} = \sqrt{\left(\frac{\delta R}{R}\right)^2 + \gamma^4 \left(\frac{\delta \beta}{\beta}\right)^2}$$

For a fixed $E_k/n = (\gamma - 1)0.9315 \text{ GeV}$
 $\frac{\Delta M}{M} = \text{const.} \Rightarrow$ mass template scaling
Templates T_7 , T_9 and T_{10} are the three unknown mass distributions.

The "Data-Driven" approach: recipe summary

The three Be mass are similar, a linear approximation is applied. σ_a is the RMS of T_a and x_a is the median of T_a . The (linear) transformation $L_{a,b}T_a = T_b$ is the function: $x \rightarrow \frac{\sigma_a}{\sigma_b}x + \left[x_a - \frac{\sigma_a}{\sigma_b}x_b\right]$.

The same transformation but applied to a different, template $L_{a,b}T_c = T_d$ is:

$$\sigma_d = \sigma_c \frac{\sigma_b}{\sigma_a} \text{ and } x_d = x_b + (x_c - x_a) \frac{\sigma_b}{\sigma_a}$$

The known (measured) data distribution is $D(x)$, thus this system must be solved:

$$D(x) = ^7\text{Be}T_7 + ^9\text{Be}T_9 + ^{10}\text{Be}T_{10}$$

$$L_{7,9}D(x) = ^7\text{Be}T_9 + ^9\text{Be}L_{7,9}T_9 + ^{10}\text{Be}L_{7,9}T_{10}$$

$$L_{7,10}D(x) = ^7\text{Be}T_{10} + ^9\text{Be}L_{7,10}T_9 + ^{10}\text{Be}L_{7,10}T_{10}$$

The ^7Be template can be written as:

$$T_7 = \frac{1}{\tau_{Be}} \left[D - \frac{^9\text{Be}}{\tau_{Be}} L_{7,9}D - \frac{^{10}\text{Be}}{\tau_{Be}} L_{7,10}D \right] + \left(\frac{^9\text{Be}}{\tau_{Be}} \right)^2 T_{G1} + \frac{^9\text{Be}}{\tau_{Be}} \frac{^{10}\text{Be}}{\tau_{Be}} (T_{G2} + T_{G3}) + \left(\frac{^{10}\text{Be}}{\tau_{Be}} \right)^2 T_{G4}$$

the last four terms, are defined by:




$$T_{G1} = L_{7,9}T_9 = L_{7,x_{G1}}T_7 \quad T_{G2} = L_{7,9}T_{10} = L_{7,x_{G2}}T_7$$

$$T_{G3} = L_{7,10}T_9 = L_{7,x_{G3}}T_7 \quad T_{G4} = L_{7,10}T_{10} = L_{7,x_{G4}}T_7$$

T_7 can be iteratively evaluated for each fixed $^7\text{Be} > ^9\text{Be} > ^{10}\text{Be}$ configuration

T_9 and T_{10} are obtained by scaling T_7 and a χ^2 is evaluated. Three un-physical $\chi^2 = 0$ solutions are $^n\text{Be}/\text{Be}=1$. Use of the statistical bootstrap is suggested for confidence intervals of physical solution.

"Data-Driven" applied to PAMELA Be events: a new $^{10}\text{Be}/^9\text{Be}$ measurement



Data-Driven approach allows a new measurement in 0.25-0.85 GeV/n. Green shaded area is a (cautious) systematic error.

First experimental hint for time dilation effect in $^{10}\text{Be}/^9\text{Be}$.

Adopting a minimal model: $^{10}\text{Be}/^9\text{Be} = Ae^{-\frac{T}{\gamma\tau}}$ (known $\tau = 2\text{My}$)

$A = 0.27 \pm 0.13$ and $T = 1.9 \pm 1.1 \text{ My}$ (dominated by PAMELA data)

Data-Driven approach allows an independent test of Montecarlo systematics. Next step is the measurement using AMS-02 data.