



The all-particle cosmic ray energy spectrum measured with HAWC




Jorge Antonio Morales Soto*,
Juan Carlos Arteaga Velázquez
for the HAWC collaboration.

Instituto de Física y Matemáticas - Universidad Michoacana

*speaker




International Cosmic Ray Conference, Berlin 2021 (ONLINE)
July, 2021.



OUTLINE


1. Introduction.
2. The HAWC Observatory.
3. Analysis and results.
4. Conclusions.

Introduction



1.1 ENERGY SPECTRUM OF COSMIC RAYS


HAWC's previous result: measurement of the all-particle energy spectrum from 10 to 500 TeV with 8 months of data [1].



Our main goals are:



- To extend this study up to 10^{15} eV with HAWC using improved statistics.
- Reduce the systematic effects using simulations that make a better description of the detector.

The HAWC Observatory



2.1 HAWC

- HAWC has as scientific objectives: to extend astrophysical measurements of gamma rays up to 100 TeV, as well as to study cosmic rays between 100 GeV and 1 PeV.
- Located between Pico de Orizaba and Sierra Negra volcanoes in Puebla, México.
- 4100 m a.s.l.
- Area of 22000 m² (62% physical coverage).
- 300 Water Cherenkov detectors.
- 1200 photomultipliers.




2.2 SIMULATIONS

- 1.3×10^7 showers were simulated with [Corsika](#) (v7.4) [2].
- Hadronic interaction models: [FLUKA](#) [3] ($E < 80$ GeV) and [QGSJet-II-04](#) [4] ($E \geq 80$ GeV).
- The interactions between secondary particles and HAWC's detectors were simulated with [GEANT4](#) [5].
- Simulated nuclei: [H](#), [He](#), [C](#), [O](#), [Ne](#), [Mg](#), [Si](#), [Fe](#). Spectra were weighted according to fits to [AMS-2](#) [6,7], [CREAM I - II](#) [8,9], and [PAMELA](#) [10]. Details of the nominal composition model are given in [1].
- $E = 5$ GeV - 3 PeV.
- Shower cores are [homogeneously distributed](#) over a circular area with 1000 m of radius centered at HAWC, with a simulated zenith angle between 0° and 70° , azimuthally symmetric. Also, they are weighted to a $\sin\theta\cos\theta$ arrival distribution.

2.3 DATA SELECTION


- Some **quality cuts** were applied to HAWC's simulated and measured data to **diminish the systematic effects** in the **energy resolution**, the **core position** and the **arrival direction**.
- Selected events:
 - Zenith angle $\theta < 35^\circ$,
 - activated at least 60 channels in a radius of 40 m from the shower core,
 - fell inside HAWC's area,
 - registered signal in, at least, 75 channels from a total of 1200,
 - and activated more than 30% of the available channels.

Analysis and results



3.1 HAWC'S MEASURED DATA

- Data from January 1st, 2018 to December 31st, 2019 were selected for this work.
- Only air showers within $E = 10^{3.5} - 10^6$ GeV were employed.



$N(E^R)$: Measured energy distribution after quality cuts

Total effective time	#events before cuts	#events after cuts
703 days	1.3638×10^{12}	1.5052×10^{10}

3.2 ENERGY SPECTRUM ESTIMATION

From $N(E^R)$ we get the unfolded energy distribution $N(E)$

How? Iterative procedure, [Bayesian Unfolding](#) [11-13]

1) $P(E_j^R | E_i)$ Response Matrix
(calculated from MC data)

2) $P(E_i | E_j^R) = \frac{P(E_j^R | E_i)P_0(E_i)}{\sum_l^{n_c} P(E_j^R | E_l)P_0(E_l)}$ Bayes formula

3) $N(E_i) = \sum_{j=1}^{n_E} P(E_i | E_j^R)N(E_j^R) = \sum_{j=1}^{n_E} M_{ij}N(E_j^R)$ True event distribution


4) $P(E_i) \equiv \frac{N(E_i)}{\sum_{i=1}^{n_c} N(E_i)} = \frac{N(E_i)}{N_{true}}$ Final probability

5) $WMSE = \frac{1}{n} \sum_{i=1}^n \frac{\bar{\sigma}_{stat,i}^2 + \bar{\delta}_{bias,i}^2}{N(E_i)}$ Weighted mean squared error
(The minimum is employed as a stopping criteria for the iteration depth)

3.2 ENERGY SPECTRUM ESTIMATION


Inputs from MC data

Response Matrix



HAWC's response becomes linear for
 $E > 10^4 \text{ GeV}$


Effective Area



Maximum trigger and reconstruction efficiency for
 $E > 10^{4.5} \text{ GeV}$

$$A_{\text{eff}}(E) = A_{\text{thrown}} \cdot \epsilon(E)$$

3.2 ENERGY SPECTRUM ESTIMATION




All-particle cosmic ray energy spectrum measured with HAWC

$$\Phi(E) = \frac{N(E)}{\Delta E \Delta t \Delta \Omega A_{eff}}$$

Contributions to the systematic error band:

1. PMT charge,
2. PMT efficiency,
3. PMT late light,
4. PMT threshold,
5. composition model (Poligonato model [14], the GSF model [15], and two models derived from fits to measurements from ATIC-2 [16] and JACEE [17]),
6. effective area,
7. seed and smoothing in unfolding,
8. unfolding technique (checked with reduced cross-entropy [18]),
9. bin size.

3.3 UNCERTAINTIES ON THE FLUX



Statistical relative error @ 10^5 GeV:

This work: +1% -0.9%


HAWC 2017 [1]: $\ll 1\%$

Systematic relative error @ 10^5 GeV:

This work: +8.2% -3.6%

HAWC 2017 [1]: +26.4% -24.7%

3.3 ALL-PARTICLE COSMIC RAY ENERGY SPECTRUM



The all-particle cosmic ray energy spectrum obtained in this work compared with the results from direct and indirect cosmic ray experiments [14-22].

3.3 FIT OF THE SPECTRUM

$$\Phi(E) = \Phi_0 E^{\gamma_1} \quad \text{Power Law}$$

$$\Phi_0 = 10^{4.38 \pm 0.01} m^{-2} s^{-1} sr^{-1} GeV^{-1}; \quad \gamma_1 = -2.632 \pm 0.001$$

$$\chi_0^2 = 492.23, \quad NDOF = 18.$$

$$\Phi(E) = \Phi_0 E^{\gamma_1} \left[1 + \left(\frac{E}{E_0} \right)^\epsilon \right]^{(\gamma_2 - \gamma_1)/\epsilon} \quad \text{Broken Power Law}$$

$$\Phi_0 = 10^{4.01 \pm 0.05} m^{-2} s^{-1} sr^{-1} GeV^{-1}$$


$$\epsilon = 3.04 \pm 1.51$$

$$\gamma_1 = -2.54 \pm 0.01$$

$$E_0 = (69.1 \pm 7.5) \text{ TeV}$$

$$\gamma_2 = -2.72 \pm 0.01$$

$$\chi_1^2 = 0.61, \quad NDOF = 15.$$



p-value for T_{obs} is

$$p < 2 \times 10^{-6}$$

4.6σ


$$TS = -\Delta\chi^2 = -(\chi_1^2 - \chi_0^2)$$

$$TS_{obs} = 491.62$$

By generating toy MC spectra with correlated data points using our covariance matrix and the result of the fit with the power-law model [23], it was found:



Conclusions



4 CONCLUSIONS



- We have extended the measurements of the total energy spectrum of cosmic rays with HAWC up to 1 PeV using a data set with high-statistics. When comparing the systematic uncertainties between this result and the result from HAWC in 2017 [1], the systematic uncertainty was reduced at $E = 10^5$ GeV. The statistical error reported in [1] only considers the size of the experimental data.
- The spectrum from this work is in agreement with the measurements from HAWC obtained in 2017 [1] and the results from NUCLEON [18].
- We also confirm the observation of a knee-like structure in the total spectrum of cosmic rays in the TeV energy regime. The position of the bump was found at $E = (69.1 \pm 7.5)$ TeV in this study.
- In addition to the measurements of NUCLEON [18], HAWC's result on the all-particle energy spectrum offer a bridge between direct and indirect measurements of the cosmic ray spectrum in the 10 TeV - 1 PeV range.

Project supported by:

Proyecto Conacyt A1-S-46288 and Coordinación de la Investigación Científica de la UMSNH.

BIBLIOGRAPHY

- [1] R. Alfaro et al., *PRD* 96 (2017) 122001.
- [2] D. Heck et al., Report No. FZKA 6019, Forschungszentrum Karlsruhe-Wissenschaftliche Berichte (1998).
- [3] A. Ferrari, et al., CERN-2005-10 (2005), INFN/TC 05/11, SLAC-R-773; G. Battistoni et al., AIP Conf. Proc. 896 (2007) 31.
- [4] S. Ostapchenko , *Phys. Rev. D* 83 (2011) 014018.
- [5] S. Agostinelli et al., *NIMA* 506 (2003) 250.
- [6] M. Aguilar et al. (AMS Collaboration), *Phys. Rev. Lett.* 114, 171103 (2015).
- [7] M. Aguilar et al. (AMS Collaboration), *Phys. Rev. Lett.* 115, 211101 (2015).
- [8] H. S. Ahn et al. (CREAM Collaboration), *Astrophys. J.* 707, 593 (2009).
- [9] Y. S. Yoon et al. (CREAM Collaboration), *Astrophys. J.* 728, 122 (2011).
- [10] O. Adriani et al. (PAMELA Collaboration), *Science* 332, 69 (2011).
- [11] D'Agostini, G. (1995). *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 362(2-3), 487-498.
- [12] Richardson, W. H. (1972). *JoSA*, 62(1), 55-59.

BIBLIOGRAPHY

[13] Lucy, L. B. (1974). *The astronomical journal*, 79, 745.

[14] . R. Hörandel, *Astrop. Phys.* 19(2) (2003) 193.

[15] . Dembinski, R. Engel, A. Fedynitch, et al. ArXiv, preprint arXiv:1711.11432, 2017.

[16] . D. Panov et al., *Bull. Russ. Acad. Sci. Phys.* 71 (2007) 494.

[17] Takahashi et al., *Nucl. Phys. B (Proc. Suppl.)* 60B (1998) 83.

[18] M. Schmelling, *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment* vol. 340, no. 2, pp. 400–412, 1994.

[19] A. D. Panov, et al., *Bulletin of the Russian Academy of Sciences: Physics* 73.5 (2009): 564-567.

[20] R. Koirala, T. K. Gaisser, et al., *POS (ICRC2019)* 318.

[21] W. D. Apel, et al., *Astroparticle Physics*, 47 (2013), 54-66.

[22] Tea Antoni, et al., *Astroparticle Physics* 24 (2005) 1–25.

[23] V. Grebenyuk, et al., *Advances in Space Research* 64.12 (2019): 2546-2558.

[24] Prosin, V. V., Berezhnev, et al., (2014). *Nuclear Instruments and Methods in Physics Research Section A: Accelerators, Spectrometers, Detectors and Associated Equipment*, 756, 94-101.

BIBLIOGRAPHY

[25] Montini, P., & Mari, S. M. (2016). arXiv preprint arXiv:1608.01389.

[26] Amenomori, M., Bi, X. J., Chen, D., Cui, S. W., Ding, L. K., Ding, X. H., ... & Tibet ASy Collaboration. (2008). The Astrophysical Journal, 678(2), 1165.

[27] Prosin, V. V., Astapov, I. I., Bezyazeev, P. A., Boreyko, V., Borodin, A. N., Brueckner, M., ... & Yashin, I. I. (2019). Energy Spectrum of Primary Cosmic Rays, According to TUNKA-133 and TAIGA-HiSCORE EAS Cherenkov Light Data. Bulletin of the Russian Academy of Sciences: Physics, 83(8), 1016-1019.

[28] P. A. Zyla et al. (Particle Data Group), Prog. Theor. Exp. Phys. 2020, 083C01 (2020) and 2021 update